

CORE EXCITED FANO-RESONANCES IN EXOTIC NUCLEI

S.E.A.Orrigo^{a,b,*}, H.Lenske^c, F.Cappuzzello^a, A.Cunsolo^{a,b}, A.Foti^{b,d}, A.Lazzaro^a,
C.Nociforo^a, J.S.Winfield^a.

a) INFN Laboratori Nazionali del Sud, Catania, Italy

b) Dipartimento di Fisica e Astronomia, Università di Catania, Catania, Italy

c) Institut für Theoretische Physik, Universität Giessen, Giessen, Germany

d) INFN Sezione di Catania, Catania, Italy

*) e-mail: orrigo@lns.infn.it

Over the last few years various unexpected phenomena have been observed in the study of light exotic nuclei, indicating a shift away from mean-field dynamics towards a new type of correlation dynamics [1]. Here, we consider a particular class of states above the neutron emission threshold, the so-called Fano-resonances, which are investigated as a new continuum excitation mode in exotic nuclei [2]. By theoretical model calculations we show that the coupling of a single particle elastic channel to closed core-excited channels leads to sharp resonances in the low-energy continuum of the $[n + (A - 1)]$ system [2]. These states are known as Bound States Embedded in the Continuum (BSEC) [3, 4, 5], which, for example, we have observed in ^{11}Be [6, 7] and ^{15}C [8, 9].

A signature for such BSEC are characteristic interference effects leading to asymmetric line shapes. We have observed a such interference in the ^{15}C spectra [8, 9]. Fano interference [10] consists of the quantum-mechanical interaction between discrete and continuous configurations, leading to characteristically asymmetric peaks in the spectra. The first model for auto-ionizing atomic states was developed by Fano in the 1960's [10]. Following the Quasiparticle-Core Coupling model [11] we consider the coupling of 1-QP (one-quasiparticle) and 3-QP components and find a number of long-living resonance structures close to the particle threshold [2]. The approach described here also extends the QPC model into the continuum. Results of the theoretical calculations for the ^{15}C nucleus are compared with the experimental data. The experimentally observed spectral distribution (both the narrow resonances and the interference pattern) are reproduced in a qualitative way, showing an agreement with a BSEC interpretation [2].

References

- [1] Lenske H., J. Phys. G: Nucl. Part. Phys. 24 (1998) 1429, and refs. therein.
- [2] Orrigo S.E.A., Lenske H. et al., Phys. Lett. B 633 (2006) 469.
- [3] Mahaux C. and Weidenmüller H.A., Shell Model Approach to Nuclear React. (1969).
- [4] Baur G. and Lenske H., Nucl. Phys. A 282 (1977) 201.
- [5] Fuchs H. et al., Nucl. Phys. A 343 (1980) 133.
- [6] Cappuzzello F. et al., Phys. Lett. B 516 (2001) 21.
- [7] Cappuzzello F. et al., Nucl. Phys. A 739 (2004) 30.
- [8] Orrigo S. E. A. et al., Proceedings 10th Int. Conf. Varenna (2003) edited by E. Gadioli, p. 147; Orrigo S. E. A., Ph.D. Thesis, University of Catania (2004).
- [9] Cappuzzello F., Orrigo S. E. A. et al., Europhys. Lett. 65 (2004) 766.
- [10] Fano U., Phys. Rev. 124 (1961) 1866.
- [11] Lenske H., Keil C. M. and Tsoneva N., J. Progr. Part. Nucl. Phys. 53 (2004) 153; Nociforo, C. and Lenske H., in preparation.