




AGATA workshop
March 23, 2004

Structure of Λ Hypernuclei Studied by γ -Ray Spectroscopy

Dept. of Physics, Tohoku University
H. Tamura

Contents

1. Introduction

Motivations and “Hyperball”

2. Structure of Hypernuclei

“Hypernuclear fine structure” and ΛN force

3. Future Plans of γ spectroscopy

“J-PARC” and “Hyperball-J” / “AGATA”

4. Summary

Hyperball collaboration (1998, 2001,2002)

E930, E509, E518

Tohoku Univ.	<u>H. Tamura</u> , Y. Fujii, O. Hashimoto, M. Kameoka, A. Matsumura, Y. Miura, T.Miyoshi, K. Mizunuma, S.N.Nakamura, H. Nomura, Y. Okayasu, T. Takahashi, M. Ukai, H. Yamauchi
Kyoto Univ.	H. Akikawa, Y.Fukao, K. Imai, K. Miwa, M.Niiyama, S.Ota, H. Takahashi, S.Terashima, M. Togawa
KEK Toyoda	Y. Kakiguchi, T. Maruta, T. Nagae, H. Noumi, H. Outa, Y. Sato, M. Sekimoto, A.
BNL	D.E. Alburger, R.E. Chrien, H. Hotchi, M. May, P. Pile, A. Rusek, R. Sutter
Osaka Univ.	S. Ajimura, T. Hayakawa, T. Kishimoto, S. Minami
CMU	G.B. Franklin, B.P. Quinn
GSI	A. Banu, T. Saitoh
Hampton Univ.	L.Tang, L. Yuan
Osaka EC Univ.	T. Fukuda, P.K.Saha,
RIKEN	<u>K. Tanida</u>
CIAE	S.H. Zhou
Univ. Freiburg	J. Franz
Science U. Tokyo	W. Imoto
ITEP	A. Krutenkova

Why Hypernuclei?

-- Extending Our Knowledge on Matter

$N_u \sim N_d \sim N_s$





$S = -\infty$

Strangeness in neutron stars ($\rho > 3 - 4 \rho_0$)

Strange hadronic matter ($A \rightarrow \infty$)

$p, n, \Lambda, \Xi^0, \Xi^-$

↑ higher density



Strangeness

$S = -2$


$S = -1$

$\Lambda\Lambda, \Xi$ hypernuclei

Λ, Σ hypernuclei

Proton-rich nuclei

Neutron-rich nuclei



non-strange nuclei

neutron number

3-dimensional nuclear chart


direct reactions

- $(K^-, \pi^-), (\pi^+, K^+) \quad n \rightarrow \Lambda$
- $(K^-, \pi^0), (\pi^-, K^0), (e, e'K^+) \quad p \rightarrow \Lambda$
- $(K^-, \pi^+) \quad pp \rightarrow \Lambda n$
- $(\pi^+, K^0) \quad nn \rightarrow \Lambda p$

Experimentally known bound hypernuclei (B_Λ known)

Must be bound

Critically bound or unbound ?



Λ Hypernuclear Chart

Motivations of hypernuclear γ spectroscopy

■ Baryon-baryon interactions


Λ hypernuclear structure $\rightleftharpoons$ ΛN int. (meson or quark models)

 c.f. nuclear structure $\longleftarrow$ NN int. (phenomenologically known)

Unified understanding of B-B interactions


■ Impurity effects

*Change of size and shape (shrinkage),
New symmetries, Change of collective motions,...*




■ Medium effects probed by hyperons free from Pauli

Λ -spin-flip $B(M1) \rightarrow g_{\Lambda}$




Hotchi et al., PRC 64 (2001) 044302

$^{89}\text{Y} (\pi^+, K^+) ^{89}_{\Lambda}\text{Y}$



Λ Single particle states
→ potential depth


$U_{\Lambda} = -30 \text{ MeV}$
 $V_{\Lambda N} < V_{NN}$



YN forces from hypernuclear structure


ΛN spin-spin, spin-orbit, tensor forces ?
 $\Lambda N \leftrightarrow \Sigma N$ coupling force?

=> precise data necessary



γ spectroscopy and Λ N spin-dependent interactions

- Low-lying levels of Λ hypernucleus



- 2-body Λ N effective interaction

Dalitz and Gal, Ann. Phys. 116 (1978) 167
 Millener et al., Phys. Rev. C31 (1985) 499

$$V_{\Lambda N}^{\text{eff}} = V_0(r) + \underbrace{V_\sigma(r)}_{\Delta} \vec{s}_\Lambda \vec{s}_N + \underbrace{V_\Lambda(r)}_{S_\Lambda} \vec{l}_{\Lambda N} \vec{s}_\Lambda + \underbrace{V_N(r)}_{S_N} \vec{l}_{\Lambda N} \vec{s}_N + \underbrace{V_T(r)}_T \vec{s}_{12}$$


p-shell : 4 radial integrals for $p_N s_\Lambda$ w.f.

Hyperball

(Tohoku/ Kyoto/ KEK, 1998)

- Large acceptance for small hypernuclear γ yields
Ge (r.e. 60%) x 14
 $\Omega \sim 15\%$
 $\varepsilon_{\text{peak}} \sim 3\%$ at 1 MeV
- High-rate electronics for huge background
1 TeV/sec, 100 kHz
- BGO counters for π^0 and Compton suppression

Resolution of hypernuclear spectroscopy
1 MeV $\rightarrow$ 2 keV FWHM





BNL-AGS: 24 GeV p, $2 \times 10^{13}/\text{spill}$
 $\rightarrow 0.9 \text{ GeV}/c \text{ K}^-$, $2 \times 10^5/\text{spill}$

Hypernuclear production by $\text{K}^- n \rightarrow \Lambda \pi^-$
 $d\sigma/d\Omega \sim 1\text{--}100 \mu\text{b}/\text{sr}$

E930 Setup

$A_Z (\text{K}^-, \pi^- \gamma) A_{\Lambda} Z$




side view

Sept.-Nov., 2001

-  drift chambers
-  Cerenkov counters
-  plastic counters


Present status

${}^7\text{Li} (\pi^+, K^+\gamma)$ KEK E419




PRL 84 (2000) 5963
PRL 86 (2001) 1982
PLB 579 (2004) 258

${}^9\text{Be} (K^-, \pi^-\gamma)$ BNL E930-1




PRL 88 (2002) 082501

${}^{13}\text{C} (K^-, \pi^-\gamma)$ BNL E929 (Nal)




PRL 86 (2001) 4255

${}^{16}\text{O} (K^-, \pi^-\gamma)$ BNL E930-2




prelim.

${}^{10}\text{B} (K^-, \pi^-\gamma)$ BNL E930-2





${}^7\text{Li} (\pi^+, K^+\gamma)$ KEK E518




-> "Table of hyper-isotopes"

${}^7_{\Lambda}\text{Li}$ Best-studied Hypernucleus



BNL E930('01)



$\gamma\gamma$ coincidence after loose event selection




First $\gamma\gamma$ coincidence for hypernuclei



${}^7_{\Lambda}\text{Li}$ Best-studied Hypernucleus

E419: ${}^7\text{Li} (\pi^+, K^+) {}^7_{\Lambda}\text{Li}$

E930-2: ${}^{10}\text{B} (K^-, \pi^-) {}^{10}_{\Lambda}\text{B}^* \rightarrow {}^7_{\Lambda}\text{Li}^*$




Non-spin-flip in weak decay $\Lambda \rightarrow N \pi$

$${}^7_{\Lambda}\text{Li} (1/2^+) \rightarrow {}^7_{\Lambda}\text{Be}^* (1/2^-) + \pi^-$$

$${}^7_{\Lambda}\text{Li} (3/2^+) \rightarrow {}^7_{\Lambda}\text{Be}^* (3/2^-) + \pi^-$$


g.s.spin = 1/2




Sasao et al., PLB 579 (2004) 258

${}^9_{\Lambda}\text{Be}$ ΛN spin-orbit force

BNL E930('98)



${}^9\text{Be} (K^-, \pi^- \gamma) {}^9_{\Lambda}\text{Be} \quad 43 \pm 5 \text{ keV}$



- Hiyama et al: *PRL* 85 (2000) 270
 $E(3/2^+) - E(5/2^+)$

Meson exch. 0.08 - 0.20 MeV

Quark model 0.035 - 0.040 MeV

-> Favors quark models

- Millener:
 $E(3/2^+) - E(5/2^+)$
 $= -0.035 \Delta - 2.465 S_{\Lambda} + 0.936 T$


$S_{\Lambda} = -0.01 \text{ MeV}$

**Consistent with
very small LS splitting
in ${}^{13}_{\Lambda}\text{C}$**

$^{16}_{\Lambda}\text{O}$ and ΛN tensor force

ΛN tensor force:

How is the nuclear tensor force without pion ?




Tensor force really very small?

Explained by meson exchange models?

$$p_{3/2}^{-1}s_{1/2} (^{12}_{\Lambda}\text{C}) : \Delta E = \frac{2}{3} \Delta + \frac{4}{3} S_{\Lambda} - \frac{8}{5} T$$




$$p_{1/2}^{-1}s_{1/2} (^{16}_{\Lambda}\text{O}) : \Delta E = -\frac{1}{3} \Delta + \frac{4}{3} S_{\Lambda} + 8 T$$



γ spectrum of $^{16}_{\Lambda}\text{O}$



$^{16}\text{O} (\text{K}^-, \pi^- \gamma) ^{16}_{\Lambda}\text{O}$

$^{16}\text{O} (\text{K}^-, \pi^-) ^{16}_{\Lambda}\text{O}$
mass spectrum



Fine structure of $^{16}_{\Lambda}\text{O}$ and ΛN tensor force

Simulated peak shape
for a fast transition




$$\Delta E = 26.1 \pm 2.0 \text{ keV (prelim.)}$$

$$E(1^-) - E(0^-) = -0.39 \Delta + 1.4 S_{\Lambda} - 0.005 S_N + 7.8 T + \Lambda \Sigma$$

(Millener, 2001)

$$\leftarrow \Delta = 0.46 \text{ MeV}, \quad S_{\Lambda} = -0.01 \text{ MeV}$$

$$\rightarrow T = 26 - 32 \text{ keV} \quad \text{First info. on } T$$




	ND	NF	NSC89	NSC97f
T	18	33	36	54

(keV)

Some of
Meson Exchange model
predictions agree


Hypernuclear Fine Structure



All the spin-dependent force parameters
determined.

J-PARC (Japan Proton Accelerator Research Complex)

By KEK+JEARI, completed in 2008 at Tokai, Japan




J-PARC under construction

Linac

3 GeV


Pacific Ocean

December, 2003




New Hyperball for J-PARC

Hyperball-J



- $\epsilon \sim 10\%$ at 1 MeV (x4 of Hyperball)
 - Rate limit: 500 kHz per each Ge (x5)
 - K beam: $\sim$ x100
- => Yield: x400 for single γ
x8000 for $\gamma\gamma$



Plans of γ spectroscopy at J-PARC (+ Hyperball-J)

(1) Complete study of light ($A < 30$) hypernuclei

(2) Systematic study of medium and heavy hypernuclei

$(K^-, \pi^- \gamma)$ spin-flip/ no-flip productions

ΛN force ($\Lambda N - \Sigma N$, p-wave,...)

$B(E2) \rightarrow$ shrinkage, New symmetries

(3) n-rich/ p-rich/ mirror hypernuclei

$(K^-, \gamma \gamma)$ in-beam method

Charge sym. break in ΛN force

$(K^-, \pi^0 \gamma)$ mirror hypernuclei

Shrinkage of n-halo

(4) Spin-flip $B(M1)$

$(K^-, \pi^- \gamma), (\pi^+, K^+ \gamma)$ Doppler shift atten.


μ_Λ in nuclei (ρ -dependence)

$(K^-, \pi^- \gamma p)$ γ -weak coincidence

(5) Double strangeness (Ξ^- atom X rays and $\Lambda\Lambda$ -hypernuclei)

$(K^-, K^+ \gamma)$




$\Xi N, \Lambda\Lambda$ interactions




Light hypernuclei

--- example of $^{12}_{\Lambda}\text{C}$

Simulation: K1.1, 10 g/cm^2 , 120 hours, Hyperball-J



$\theta_{\pi\gamma}$ angular correlation -> spin assignment



--- Disappearance
of n-halo in ${}^7_{\Lambda}\text{He}$

Energy level diagram showing the transition from ${}^6\text{He}$ to ${}^7_{\Lambda}\text{He}$.

Y-axis: Energy E (MeV).

${}^6\text{He}$ levels:

- 2^+ at 0 MeV (crossed out).
- 0^+ at -0.69 MeV.

${}^7_{\Lambda}\text{He}$ levels:



- $5/2^+$ at -4.43 MeV.
- $3/2^+$ at -4.43 MeV.
- $1/2^+$ at -6.12 MeV.

Transitions:

- $B_{\Lambda} = 5.44$ MeV (dashed arrow from ${}^6\text{He}$ 0^+ to ${}^7_{\Lambda}\text{He}$ $1/2^+$).
- $E2$ (red arrow from ${}^7_{\Lambda}\text{He}$ $3/2^+$ to $1/2^+$).

Drastic $B(E2)$ change

0.58	$B(E2) \text{ e}^2\text{fm}^4$	0.068
		0.059



AGATA(500kHz) at J-PARC?

-> $\varepsilon_\gamma \times 5$, $\varepsilon_{\gamma\gamma} \times 25$ of Hyperball-J

New experimental possibilities open.

- Impurity effect of Λ :
 Change of deformation, collective motions, ..
 Shrinkage by $\Lambda\Lambda$ -> Density of strange nuclear matter
- Various n-rich hypernuclei such as $^{11}_{\Lambda}\text{Li}$, $^{12}_{\Lambda}\text{Li}$
 $(K^-, \pi^+) \sim \text{nb/sr}$ $^{11}\text{B} (K^-, \pi^+) \text{ } ^{11}_{\Lambda}\text{Li}$ (Λ makes ^{10}Li bound)
 K^- -in-beam: $(K^-, \gamma\gamma)$ $^{12}_{\Lambda}\text{Li}$
- Systematic spectroscopy of various $\Lambda\Lambda$ hypernuclei
 $Z (K^-, K^+) \Xi [Z-2] \rightarrow {}_{\Lambda\Lambda}Z' + \dots$

Summary


- **Hypernuclear γ spectroscopy with Hyperball in progress**
-> ΛN interaction, shrinking effect from $B(E2)$
First success of $\gamma\gamma$ coincidence
- **${}^7_{\Lambda}\text{Li}$, ${}^9_{\Lambda}\text{Be}$, ${}^{16}_{\Lambda}\text{O}$: Hypernuclear fine structure**
-> **All the ΛN spin-dependent force parameters**
 $(\Delta, S_{\Lambda}, S_N, T)$ determined.
LS (S_{Λ}, S_N) quark picture favored
Tensor (T) meson exchange OK
- **J-PARC high-intensity K beam + Hyperball-J (or AGATA)**
-> **“Table of Hyper-isotopes” incl. heavy, n-rich,.. hypernuclei**
AGATA will further extend possibilities.

Λ -Spin-flip B(M1) $B(M1) \sim |\langle \Phi$

μ_Λ in nucleus $\rightarrow$ medium effect of baryons



$$\frac{e\hbar}{2mc} ?$$



$$\begin{aligned} B(M1) &\propto |\langle \Phi_f | \mu^z | \Phi_i \rangle|^2 \\ &= |\langle \phi_{\Lambda \uparrow} \phi_c | g_c J_c^z + g_\Lambda J_\Lambda^z | \phi_{\Lambda \downarrow} \phi_c \rangle|^2 \\ &\propto (g_c - g_\Lambda)^2 \end{aligned}$$


■ Doppler shift attenuation method [same as $B(E2)$, *established*]
 $(\pi^+, K^+ \gamma)$, $(K^-, \pi^- \gamma)$ for light hypernuclei

■ γ -weak coincidence method [*new, only possible at J-PARC*]
 $(K^-, \pi^- p \gamma)$ for heavy hypernuclei

Accuracy $\Delta g_\Lambda < 5\%$ achievable

$\rightarrow$ ρ dependence of g_Λ

Setup at K1.1



■ Beamline: K1.1 (K1.1BR)

K^- at 1.1, 0.8 GeV/c

4.2×10^7 K-/spill

$K/\pi > 1$

π^+ at 1.05 GeV/c

■ Spectrometer: SPESII (existing)

$\Delta p/p < 2$ MeV (FWHM)

$\Omega \sim 20$ msr


■ Hyperball-J (2007?)

■ New CDS

(K⁻,K⁺) experiments with K1.8 + SKS

- Ξ hypernuclear spectroscopy
- $\Lambda\Lambda$ hypernuclei
+ CDS for decay π 's
hybrid emulsion
weak decays
 γ rays

↓
 ΞN , $\Lambda\Lambda$, ΞN - $\Lambda\Lambda$
interactions




1.8 GeV/c K⁻ beam
1.0x10⁷/pulse

(1-b) Light hypernuclei


Impurity effect

--- example of $^{20}_{\Lambda}\text{Ne}$: change of cluster structure



(a) Millener et al.

(b) Sakuda - Bando



Strangeness Nuclear Physics

Experiments at J-PARC

Day-1 experiments

- K1.8** ■ Ξ hypernuclear spectroscopy by (K^-, K^+)
- K1.8** ■ $\Lambda\Lambda$ hypernuclei by sequential π decay method
and by hybrid emulsion method
- K1.1** ■ Precision γ spectroscopy of Λ and $\Lambda\Lambda$ hypernuclei, Ξ X-rays
K1.8
- HR** ■ High-resolution (π, K^+) spectroscopy for Λ , Σ , and n-rich Λ hypernuclei
- K1.1** ■ Weak decays of Λ and $\Lambda\Lambda$ hypernuclei
K1.8
- K1.1** ■ K-nucleus bound states by (K^-, N) and (K^-, π) reactions
- K1.1** ■ ΛN , ΣN , ΞN scattering experiments
K1.8

Do we understand nuclear force?

Short-range forces (=repulsive core, LS force) determine the “basic properties of nucleus” (=saturation, magic number).


Quark Cluster Model

- NN repulsive core
 Δ -N mass difference $\rightarrow$ color magnetic int.

$[\uparrow u \uparrow u \downarrow d \uparrow d \downarrow s \uparrow s \downarrow]$

H dibaryon ($SU_f(3) \times SU_f(3)$ singlet)
 = Attractive core in $\Lambda\Lambda$ - ΞN channel

- ΣN ($I=3/2, S=1$) strongly repulsive core
 $\leftarrow$ quark Pauli
- LS force $\rightarrow$ quark gluon exchange
 $\rightarrow$ Λ : very small
 Σ : as large as N



Λ spin-orbit splitting in $^{13}_{\Lambda}\text{C}$



$^{13}\text{C} (\text{K}^-, \pi^- \gamma)$

BNL E929

w/ NaI array

$p_{1/2}, p_{3/2}$ single-particle level splitting


$$E(1/2^-) - E(3/2^-) = 152 \pm 54 \pm 36 \text{ keV}$$



Ajimura et al., PRL 86 (2001) 4255

FIG. 2. γ ray spectra taken in coincidence with scattered π^- 's (upper panel) and differential cross section of $1/2^-$ and $3/2^-$ states calculated by Motoba [18] (lower panel) are shown.

How to measure Tensor term: $^{16}_{\Lambda}\text{O}$ and $^{15}_{\Lambda}\text{N}$




$p_{1/2}$ shell --- Large contribution of the tensor term


Upgrade of Hyperball

Construction of Hyperball2 in progress. Ready in 2004


- Clover Ge (r.e. >120%)
+BGO x 6 added
- Peak eff.
~ 2.5% -> 5% at 1 MeV
- Beam test of Clover Ge
(T536, June 2003) OK
- VME-based fast readout
- Improvement of preamplifier



J-PARC NP-Hall (a sample plan by Noumi)



Physics output from ${}^7_{\Lambda}\text{Li}$



shrinking effect

Predicted by
Motoba et al.,
Prog.Theor.Phys.
70 (1983) 189.

$B(E2)$

$\propto | \langle f | e r^2 Y_2 | i \rangle |^2$


$\propto R^4$

$B(E2)$
[$e^2 \text{fm}^4$]

10.9 ± 0.9

$\longrightarrow 3.6 \pm 0.5 \pm 0.5$
 $\Rightarrow 19 \pm 4\%$ Shrinkage by Λ

N- LS interaction
 $S_N \sim -0.4 \text{ MeV}$



spin-spin interaction
 $\Delta = 0.50 \text{ MeV}$